



## Problem 7. «A fixed element»

A polynomial  $f(X_1, \dots, X_n) \in \mathbb{F}_2[X_1, \dots, X_n]$  is called *reduced* if the degree of each  $X_i$  in  $f$  is at most 1. For  $0 \leq r \leq n$ , the  $r$ th order Reed – Muller code of length  $2^n$ , denoted by  $R(r, n)$ , is the  $\mathbb{F}_2$ -space of all reduced polynomials in  $X_1, \dots, X_n$  of total degree less or equal than  $r$ . We also define  $R(-1, n) = \{0\}$ .

The general linear group  $GL(n, \mathbb{F}_2)$  acts on  $R(r, n)$  naturally: Given  $A \in GL(n, \mathbb{F}_2)$  and  $f(X_1, \dots, X_n) \in R(r, n)$ ,  $Af$  is defined to be the reduced polynomial obtained from  $f((X_1, \dots, X_n)A)$  by replacing each power  $X_i^k$  ( $k \geq 2$ ) with  $X_i$ . Consequently,  $GL(n, \mathbb{F}_2)$  acts on the quotient space  $R(r, n)/R(r - 1, n)$ .

Let  $A \in GL(n, \mathbb{F}_2)$  be such that its characteristic polynomial is a primitive irreducible polynomial over  $\mathbb{F}_2$ . Prove that the only element in  $R(r, n)/R(r - 1, n)$  fixed by the action of  $A$  is 0.

